
Rubus phoenicolasius

Distribution






Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **1–2**
Life form: **nanophanerophyte**
Life strategy: **C – competitor**
Life strategy (Pierce method based on leaf traits): **C/CSR**
Life strategy (Pierce method, C-score): **51.3 %**
Life strategy (Pierce method, S-score): **23.9 %**
Life strategy (Pierce method, R-score): **24.9 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **compound – ternate**
Stipules: **present**
Petiole: **present**
Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **June–July**
Flower colour: **pink**
Flower symmetry: **actinomorphic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **aposepalous**
Inflorescence type: **racemus**
Dicliny: **synoecious**
Pollination syndrome: **insect-pollination**



Fruit, seed and dispersal

Fruit type: **fleshy fruit – head of one-seeded drupes**
Fruit colour: **orange, red**
Reproduction type: **by seed/spores and vegetatively**
Myrmecochory: **non-myrmecochorous (b)**



Belowground organs and clonality

Root metamorphosis: **root shoot**



Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**



Taxon origin


Origin in the Czech Republic: **neophyte**
Invasion status: **casual**
Geographic origin: **Asia**
Period of introduction: **Recent Past (1950–2000)**
Introduction pathway: **intentional – ornamental, intentional – crops**



Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**



Distribution and frequency

Floristic region: **Eastern Asia**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 7
taxon.data.freq_in_quad: 8

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz