
Trifolium alpinum




Distribution



Map info

- revised records
- unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Growth form: **polycarpic perennial non-clonal herb**
Life form: **hemicryptophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **compound – ternate**
Stipules: **present**
Petiole: **present**

Flower

Flower colour: **pink**
Flower symmetry: **zygomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **synsepalous**
Inflorescence type: **capitulum**

Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**

© 2014-2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz







Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot life span (cyclicity): **monocyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **monopodial**
Primary root: **present**

Bud bank

Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **15**
Depth of the belowground bud bank (root buds exluded) [cm]: **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **15**
Depth of the belowground bud bank (root buds included) [cm]: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in the Czech Republic: **neophyte**
Invasion status: **casual**
Geographic origin: **Europe**
Year of the first record in the wild: **1919**
Period of introduction: **Late Modern Period (1800–1950)**
Introduction pathway: **intentional – nature**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 – light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**
Temperature indicator value: **2 – transition between values 1 and 3**
Moisture indicator value: **4 – transition between values 3 and 5**
Reaction indicator value: **2 – transition between values 1 and 3**
Nutrient indicator value: **2 – transition between values 1 and 3**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

does not spontaneously occur in Czech forests

Distribution and frequency

Continentality degree: **3**
Distribution range extension along the continentality gradient: **3**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 11
taxon.data.freq_in_quad: 11

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz