


# *Filipendula ulmaria* subsp. *ulmaria*

## Distribution



### Map info

● revised records

● unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **compound - interruptedly pinnate**

Stipules: **present**

Petiole: **present**

## Flower

Flower colour: **yellow-white**

Flower symmetry: **actinomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **free**

Calyx fusion: **aposepalous**

Inflorescence type: **anthella**

## Fruit, seed and dispersal

Fruit type: **dry fruit - head of achenes**

Dispersal unit (diaspore): **fruit, infructescence or its part**

Dispersal strategy: **Allium (mainly autochory)**

Myrmecochory: **non-myrmecochorous (b)**

## Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in the Czech Republic: **native**

## Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **6 - transition between values 5 and 7; rarely at less than 20% of diffuse radiation incident in an open area**

Temperature indicator value: **5x - moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas (generalist)**

Moisture indicator value: **8 - transition between values 7 and 9**

Reaction indicator value: **6x - transition between values 5 and 7 (generalist)**

Nutrient indicator value: **5 - occurring at moderately nutrient-rich sites, and less frequently at poor and rich sites**

Salinity indicator value: **1 - salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**

## Habitat and sociology

Occurrence in habitats

2 Alpine and subalpine grasslands

2B Subalpine tall-forb and tall-grass vegetation: **1 - rare occurrence**

4 Wetland and riverine herbaceous vegetation

4A Reed-beds of eutrophic still waters: **1 - rare occurrence**

4B Halophilous reed and sedge beds: **1 - rare occurrence**

4D Riverine reed vegetation: **2 - optimum**

4E Reed vegetation of brooks: **1 - rare occurrence**

4F Mesotrophic vegetation of muddy substrata: **1 - rare occurrence**

4G Tall-sedge beds: **2 - optimum**

4J River gravel banks: **1 - rare occurrence**

4K Petasites fringes of montane brooks: **2 - optimum**

4L Nitrophilous herbaceous fringes of lowland rivers: **1 - rare occurrence**

5 Vegetation of springs and mires

5B Lowland to montane soft-water springs: **1 - rare occurrence**

5C Alpine and subalpine soft-water springs: **1 - rare occurrence**

5D Calcareous fens: **2 - optimum**

5E Acidic moss-rich fens and peatland meadows: **2 - optimum**

5F Transitional mires: **1 - rare occurrence**

6 Meadows and mesic pastures

6A Mesic Arrhenatherum meadows: **1 - rare occurrence**

6B Montane mesic meadows: **1 - rare occurrence**

6C Pastures and park grasslands: **1 - rare occurrence**  
6D Alluvial meadows of lowland rivers: **2 - optimum**  
6E Wet Cirsium meadows: **3 - dominant**  
6F Intermittently wet Molinia meadows: **2 - optimum**  
7 Acidophilous grasslands  
7B Submontane Nardus grasslands: **1 - rare occurrence**  
9 Sand grasslands and rock-outcrop vegetation  
9E Acidophilous vegetation of spring therophytes and succulents: **1 - rare occurrence**  
10 Saline vegetation  
10J Saline steppes: **1 - rare occurrence**  
11 Heathlands and scrub  
11H Subalpine deciduous scrub: **1 - rare occurrence**  
11I Willow carrs: **1 - rare occurrence**  
11J Willow galleries of loamy and sandy river banks: **2 - optimum**  
11L Tall mesic and xeric shrub: **1 - rare occurrence**  
11R Scrub and pioneer woodland of forests clearings: **1 - rare occurrence**  
12 Forests  
12A Alder carrs: **2 - optimum**  
12B Alluvial forests: **2 - optimum**  
12U Plantations of broad-leaved non-native trees: **1 - rare occurrence**  
13 Anthropogenic vegetation  
13E Perennial nitrophilous herbaceous vegetation of mesic sites: **1 - rare occurrence**  
Affinity to the forest environment  
Affinity to the forest environment in Thermophyticum: **2.1 - taxon occurring both in the forest and open vegetation**  
Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.1 - taxon occurring both in the forest and open vegetation**

## Distribution and frequency

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt, montane belt**  
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 423  
taxon.data.freq\_in\_quad: 842  
Number of habitats with taxon occurrence in the Czech Republic  
Number of narrow habitats in which the taxon occurs: **33**  
Number of narrow habitats in which the taxon has its optimum: **11**  
Number of broad habitats in which the taxon occurs: **10**  
Number of broad habitats in which the taxon has its optimum: **5**

## Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**  
Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**  
Legal protection: **not protected by law**