
Gentiana cruciata subsp. *cruciata*

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.





Habitus and growth type

Height [m]: **0.12–0.45**
Growth form: **polycarpic perennial non-clonal herb**
Life form: **hemicryptophyte**
Life strategy: **CSR – competitor/stress-tolerator/ruderal**
Life strategy (Pierce method based on leaf traits): **CSR**
Life strategy (Pierce method, C-score): **38.4 %**
Life strategy (Pierce method, S-score): **32.1 %**
Life strategy (Pierce method, R-score): **29.6 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **opposite, rosulate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **both present and absent**
Leaf life span: **summer green**
Leaf anatomy: **scleromorphic, mesomorphic**

Flower

Flowering phase: **8 Clematis vitalba-Galium sylvaticum (mid-summer)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Flower colour: **blue**
Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **campanulate**
Calyx fusion: **synsepalous**
Inflorescence type: **racemus e verticillastris compositus**
Dicliny: **synoecious**
Generative reproduction type: **mixed mating**
Pollination syndrome: **insect-pollination, selfing**





Fruit, seed and dispersal

Fruit type: **dry fruit – capsule**
Fruit colour: **brown**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**





Belowground organs and clonality

Shoot metamorphosis: **pleiocorm**
Root metamorphosis: **primary storage root**
Storage organ: **pleiocorm, primary storage root**
Shoot life span (cyclicity): **monocyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **present**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **8**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **13**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **20**
Depth of the belowground bud bank (root buds exluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **8**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **13**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **20**
Depth of the belowground bud bank (root buds included) [cm]: **4**





Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**





Karyology

Chromosome number (2n): **52**



Ploidy level (x): **4**
2C genome size [Mbp]: **7430.55**
1Cx monoploid genome size [Mbp]: **1857.64**
Genomic GC content: **40.8 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **6 – transition between values 5 and 7**
Moisture indicator value: **3 – missing on damp soil**
Reaction indicator value: **8 – transition between values 7 and 9, occurring mostly in calcium-rich conditions**
Nutrient indicator value: **3 – occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Habitat and sociology

Occurrence in habitats

6 Meadows and mesic pastures

6A Mesic Arrhenatherum meadows: **1 – rare occurrence**
6C Pastures and park grasslands: **1 – rare occurrence**

8 Dry grasslands

8D Broad-leaved dry grasslands: **2 – optimum**
8E Acidophilous dry grasslands: **1 – rare occurrence**
8F Thermophilous forest fringe vegetation: **1 – rare occurrence**

11 Heathlands and scrub

11L Tall mesic and xeric shrub: **1 – rare occurrence**

12 Forests

12H Peri-Alpidic basiphilous thermophilous oak forests: **1 – rare occurrence**
12I Sub-continental thermophilous oak forests: **1 – rare occurrence**
12O Peri-Alpidic pine forests: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of associations: **TDC02 *Anthoxantho odorati-Agrostietum tenuis***

Ecological specialization indices

Ecological specialization index for all vegetation types: **6.1**
Ecological specialization index for non-forest vegetation: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Colonization ability

Index of colonization success (ICS): **1**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional**
Floristic region: **Europe, Western Asia**
Continentality degree: **6**
Distribution range extension along the continentality gradient: **5**
Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt, montane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 170
taxon.data.freq_in_quad: 269

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0 %**
Occurrence frequency in vegetation plots with a cover above 5%: **0 %**
Occurrence frequency in vegetation plots with a cover above 25%: **0 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **1.9 %**
Maximum percentage cover in vegetation plots: **3 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **9**
Number of narrow habitats in which the taxon has its optimum: **1**
Number of broad habitats in which the taxon occurs: **4**
Number of broad habitats in which the taxon has its optimum: **1**

Threats and protection

Red List 2017 (national categories): **C2b – endangered taxon, rare and declining**
Red List 2017 (IUCN categories): **EN – endangered**
Legal protection: **vulnerable taxon**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz