
# *Gentianella lutescens* subsp. *carpatica*

## Distribution






Map info

*revised records*

*unrevised records*

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







### Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **opposite, rosulate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**

### Flower

Flowering period [month]: **August–October**
Flower colour: **blue-violet**
Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **funnel-shaped**
Calyx fusion: **synsepalous**
Inflorescence type: **panicula**

### Fruit, seed and dispersal

Fruit type: **dry fruit – capsule**
Fruit colour: **brown**
Dispersal unit (diaspore): **seed**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**




## Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin


Origin in the Czech Republic: **native**




## Ecological indicator values

### Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **5 – moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**
Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **5 – indicator of moderate acidity, occurring rarely in strongly acidic as well as in neutral to alkaline conditions**
Nutrient indicator value: **3 – occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**




## Habitat and sociology

### Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**




## Threats and protection

Red List 2017 (national categories): **C1t – critically threatened taxon, declining**
Red List 2017 (IUCN categories): **CR – critically endangered**
Legal protection: **endangered taxon**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*






*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*