

Leucanthemum vulgare subsp. *vulgare*

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **0.3–0.8**
Growth form: **clonal herb**
Life form: **hemicryptophyte**
Life strategy: **C – competitor**
Life strategy (Pierce method based on leaf traits): **R/CR**
Life strategy (Pierce method, C-score): **20.9 %**
Life strategy (Pierce method, S-score): **0 %**
Life strategy (Pierce method, R-score): **79.2 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **both present and absent**
Leaf life span: **evergreen**
Leaf anatomy: **scleromorphic, mesomorphic**

Flower

Flowering period [month]: **May–October**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Flower colour: **white, yellow**
Flower symmetry: **actinomorphic, zygomorphic**
Perianth type: **calyx sometimes absent, corolla present**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **ligulate, tubular**
Calyx fusion: **synsepalous**
Inflorescence type: **anthodium solitarium**
Dicliny: **gynomonoecious**
Generative reproduction type: **mixed mating**
Pollination syndrome: **insect-pollination, selfing**



Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**
Fruit colour: **brown, black**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **myrmecochorous**



Belowground organs and clonality

Shoot metamorphosis: **stolon-like rhizome**
Storage organ: **stolon-like rhizome**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **18**
Ploidy level (x): **2**
2C genome size [Mbp]: **9802.24**
1Cx monoploid genome size [Mbp]: **4901.12**
Genomic GC content: **40 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **5x – moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas (generalist)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **6x – transition between values 5 and 7 (generalist)**
Nutrient indicator value: **5 – occurring at moderately nutrient-rich sites, and less frequently at poor and rich sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1A Calcareous cliffs: **1 – rare occurrence**
1D Mobile calcareous screes: **1 – rare occurrence**

5 Vegetation of springs and mires

5D Calcareous fens: **2 – optimum**

6 Meadows and mesic pastures

6A Mesic Arrhenatherum meadows: **2 – optimum**
6B Montane mesic meadows: **2 – optimum**
6C Pastures and park grasslands: **2 – optimum**
6D Alluvial meadows of lowland rivers: **2 – optimum**
6E Wet Cirsium meadows: **1 – rare occurrence**
6F Intermittently wet Molinia meadows: **2 – optimum**
6G Vegetation of wet disturbed soils: **1 – rare occurrence**

7 Acidophilous grasslands

7A Subalpine and montane acidophilous grasslands: **2 – optimum**
7B Submontane Nardus grasslands: **2 – optimum**

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **1 – rare occurrence**
8B Submediterranean dry grasslands on rock outcrops: **1 – rare occurrence**
8C Narrow-leaved sub-continental steppes: **1 – rare occurrence**
8D Broad-leaved dry grasslands: **2 – optimum**
8E Acidophilous dry grasslands: **1 – rare occurrence**
8F Thermophilous forest fringe vegetation: **1 – rare occurrence**

9 Sand grasslands and rock-outcrop vegetation

9B Open vegetation of acidic sands: **1 – rare occurrence**
9C Festuca grasslands on acidic sands: **1 – rare occurrence**
9E Acidophilous vegetation of spring therophytes and succulents: **1 – rare occurrence**
9F Basiphilous vegetation of spring therophytes and succulents: **1 – rare occurrence**

10 Saline vegetation

10I Inland saline meadows: **1 – rare occurrence**

11 Heathlands and scrub

11A Dry lowland to subalpine heathlands: **1 – rare occurrence**
11L Tall mesic and xeric shrub: **1 – rare occurrence**
11N Low xeric scrub: **1 – rare occurrence**

12 Forests

12C Oak-hornbeam forests: **1 – rare occurrence**
12H Peri-Alpidic basiphilous thermophilous oak forests: **1 – rare occurrence**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

12I Sub-continental thermophilous oak forests: **1 – rare occurrence**
12J Acidophilous thermophilous oak forests: **1 – rare occurrence**
12K Acidophilous oak forests: **1 – rare occurrence**
12O Peri-Alpidic pine forests: **1 – rare occurrence**
12W Pine and larch plantations: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**

Colonization ability

Index of colonization success (ICS): **7**
Index of colonization potential (ICP): **4**
Optimum successional age [years]: **24.5**

Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional**
Floristic region: **Europe, Western Siberia**
Distribution range extension along the continentality gradient: **6**
Elevational belt in the Czech Republic: **lowlands, colline belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 638
taxon.data.freq_in_quad: 1974

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **33**
Number of narrow habitats in which the taxon has its optimum: **9**
Number of broad habitats in which the taxon occurs: **9**
Number of broad habitats in which the taxon has its optimum: **4**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz