
Salix triandra subsp. *amygdalina*

Distribution



Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **present**
Petiole: **present**

Flower

Flower colour: **green**
Perianth type: **flower achlamydeous**
Inflorescence type: **amentum e floribus masculis, amentum e floribus femineis**

Fruit, seed and dispersal

Fruit type: **dry fruit – capsule**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Phragmites (mainly anemochory and hydrochory)**
Myrmecochory: **non-myrmecochorous (b)**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **5 – moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**
Moisture indicator value: **8 – transition between values 7 and 9**
Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**
Nutrient indicator value: **6 – transition between values 5 and 7**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**

Distribution and frequency

Elevational belt in the Czech Republic: **lowlands, colline belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 25
taxon.data.freq_in_quad: 29

Threats and protection

Red List 2017 (national categories): **C4b – data deficient taxon**
Red List 2017 (IUCN categories): **DD – data deficient**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz