
Sisymbrium austriacum subsp. *austriacum*

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.





Habitus and growth type

Height [m]: **0.3–0.6**
Growth form: **monocarpic perennial non-clonal herb**
Life form: **hemicryptophyte**
Life strategy: **CSR – competitor/stress-tolerator/ruderal**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – pinnately divided**
Stipules: **absent**
Petiole: **mainly present**
Leaf life span: **evergreen**
Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **May–June**
Flowering phase: **5 Sorbus aucuparia-Galium odoratum (end of mid-spring)**
Flower colour: **yellow**
Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz







Perianth fusion: **free**
Calyx fusion: **aposepalous**
Inflorescence type: **panicula**
Dicliny: **synoecious**
Generative reproduction type: **mixed mating**
Pollination syndrome: **insect-pollination, selfing**

Fruit, seed and dispersal

Fruit type: **dry fruit – siliqua**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Root metamorphosis: **primary storage root**
Storage organ: **primary storage root**
Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**
Primary root: **present**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **5**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **10**
Depth of the belowground bud bank (root buds exluded) [cm]: **3**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **5**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **10**
Depth of the belowground bud bank (root buds included) [cm]: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **neophyte**
Invasion status: **casual**
Geographic origin: **Europe, Mediterranean**
Year of the first record in the wild: **1858**
Period of introduction: **Late Modern Period (1800–1950)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**
Floristic region: **Europe**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 1
taxon.data.freq_in_quad: 1

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz