
Thymus pulegioides subsp. *montanus*

Distribution



Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **opposite**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **present**

Flower

Flower colour: **pink**
Flower symmetry: **zygomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **bilabiate**
Calyx fusion: **synsepalous**
Inflorescence type: **pseudospica e verticillastris composita**

Fruit, seed and dispersal

Fruit type: **dry fruit – cluster of four one-seeded nutlets**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **probably myrmecochorous**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **6 – transition between values 5 and 7**
Moisture indicator value: **4 – transition between values 3 and 5**
Reaction indicator value: **6x – transition between values 5 and 7 (generalist)**
Nutrient indicator value: **3 – occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**

Threats and protection

Red List 2017 (national categories): **C4b – data deficient taxon**
Red List 2017 (IUCN categories): **DD – data deficient**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz