
# *Vicia pannonica* subsp. *striata*

## Distribution






Map info

*revised records*

*unrevised records*

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.










### Habitus and growth type

Life form: **therophyte**

### Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **compound – paripinnate**
Stipules: **present**
Petiole: **present**

### Flower

Flower colour: **red-violet**
Flower symmetry: **zygomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **synsepalous**
Inflorescence type: **racemus**

### Fruit, seed and dispersal

Fruit type: **dry fruit – legume**
Dispersal unit (diaspore): **seed**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*







Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

## Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in the Czech Republic: **archaeophyte**
Invasion status: **naturalized**
Geographic origin: **Mediterranean**
Period of introduction: **Neolithic (5600–4200 BCE)**
Introduction pathway: **unintentional – agriculture**

## Ecological indicator values

### Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **7 – heat indicator, occurring in relatively warm lowlands**
Moisture indicator value: **4 – transition between values 3 and 5**
Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**
Nutrient indicator value: **6 – transition between values 5 and 7**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

## Habitat and sociology

### Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

## Distribution and frequency

Continentality degree: **7**
Distribution range extension along the continentality gradient: **4**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 139
taxon.data.freq_in_quad: 225

## Threats and protection

Red List 2017 (national categories): **C2t – endangered taxon, declining**
Red List 2017 (IUCN categories): **EN – endangered**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Legal protection: **not protected by law**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*