

Rorippa ×armoracioides

Distribution
















Habitus and growth type

Height [m]: **0.3–0.9**
Life form: **hemicryptophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Stipules: **absent**
Petiole: **both present and absent**

Flower

Flowering period [month]: **June–August**
Flower colour: **yellow**
Perianth type: **calyx and corolla**
Calyx fusion: **aposepalous**
Inflorescence type: **racemus**

Fruit, seed and dispersal

Dispersal unit (diaspore): **seed**
Dispersal strategy: **Allium (mainly autochory)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**



Taxon origin

Origin in the Czech Republic: **native**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**




Distribution and frequency

Elevational belt in the Czech Republic: **lowlands, colline belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 44
taxon.data.freq_in_quad: 54







© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz







© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz